

Product data sheet

Daudi/NF-kB Reporter (Luc) stable cell line

Catalog Number CL-1315

Storage: Liquid nitrogen

Components: 1 vial contains $\sim 2 \times 10^6$ cells in Cell freezing medium

Product description

Daudi/NF-kB Reporter (Luc) cells are derived from the human Burkitt lymphoma cell line Daudi by stably integration of a NF-kB firefly luciferase reporter construct. Daudi cells have been widely used in cancer research and drug development. Daudi/NF-kB Reporter (Luc) cells express firefly luciferase under the control of the NF-kB response elements, can be used for *in vitro* assays and *in vivo* imaging.

Cell line description

Organism: Homo sapiens (human)

Tissue: Peripheral blood

Cell Type: B lymphoblast

Morphology: Lymphoblast

Culture Properties: Suspension

Disease: Burkitt's lymphoma

Biosafety Level: 2

Medium

1. Complete culture medium: RPMI-1640, 10% fetal bovine serum (FBS)
0.5 $\mu\text{g/mL}$ of puromycin may be added to the culture medium. Puromycin should not be added until a culture has been well established from the thawed cells.
2. Freeze medium: RPMI-1640, 10% fetal bovine serum (FBS), 6% DMSO

Culture procedure

Thawing of frozen cells

1. Thaw the frozen cryovial by gentle agitation in a 37 °C water bath in 1-2 minutes.
2. Remove the cryovial from the water bath as soon as the contents are thawed, and decontaminate by wiping with 70% ethanol.
3. Transfer the thawed cell suspension to a centrifuge tube containing 10 ml of Complete culture medium, centrifuge at 500 g for 5 minutes.

4. Remove the medium by aspiration, resuspend the cells with 2 ml of the Complete culture medium by gently pipetting up and down.
5. Transfer the cells to a T-25 suspension cell culture flask.
6. Place the cells in a 37°C incubator with 5% CO₂.

Sub-culturing

Cultures can be maintained by the addition of fresh medium or replacement of medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 3 to 5×10^5 viable cells/ml. Maintain cell density between 3×10^5 and 2 to 3×10^6 viable cells/ml. Do not allow the cell density to exceed 2 to 3×10^6 cells/ml.

Renew or add fresh medium every 2-3 days.