

## Product data sheet

HEK293/GFP stable cell line

Catalog Number: CL-1121

Storage: Liquid nitrogen

Components: 1 vial contains  $\sim 2 \times 10^6$  cells in Cell freezing medium

## Product description

HEK 293/GFP cells are derived from the human immortalized embryonic kidney cell line HEK 293 by stably integration of a constitutive GFP expression construct. HEK 293 cells have been widely used in cell biology research because of their reliable growth and propensity for transfection. HEK 293/GFP cells stably express GFP, can be used for *in vitro* assays and *in vivo* imaging.




Figure 1. GFP expression in HEK293/GFP stable cell line

## Cell line description

Organism: *Homo sapiens*, human

Tissue: kidney; Embryo

Morphology: epithelial

Culture Properties: adherent

## Medium

1. Complete culture medium: DMEM with 10% fetal bovine serum (FBS)  
1  $\mu$ g/mL of puromycin may be added to the culture medium. **Puromycin should not be added until a culture has been well established from the thawed cells.**

2. Freeze medium: FBS with 6% DMSO

## **Culture procedure**

### Thawing of frozen cells

1. Thaw the frozen cryovial by gentle agitation in a 37 °C water bath in 1-2 minutes.
2. Remove the cryovial from the water bath as soon as the contents are thawed, and decontaminate by wiping with 70% ethanol.
3. Transfer the thawed cell suspension to a centrifuge tube containing 10 ml of Complete culture medium, centrifuge at 500 g for 5 minutes.
4. Remove the medium by aspiration, resuspend the cells with 10 ml of the Complete culture medium by gently pipetting up and down.
5. Transfer the cells to a 10 cm cell culture dish.
6. Place the cells in a 37°C incubator with 5% CO<sub>2</sub>.

### Sub-culturing

Volumes are given for a 10 cm cell culture dish. Increase or decrease the amount of dissociation medium needed proportionally.

1. Remove the medium by aspiration.
2. Briefly rinse the cell layer with 1xDPBS to remove all traces of serum that contains trypsin inhibitor.
3. Add 1 ml of Trypsin-EDTA (0.25%) solution to the dish and observe cells under an inverted microscope until cell layer is dispersed.
4. Add 4 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C with 5% CO<sub>2</sub>.