

Product data sheet

NCI-H929/GFP stable cell line

Catalog Number: CL-1643

Storage: Liquid nitrogen

Components: 1 vial contains $\sim 2 \times 10^6$ cells in Cell freezing medium

Product description

NCI-H929/GFP cells are derived from the human B lymphocyte NCI-H929 cell line by stably integration of a constitutive GFP stably expression construct. NCI-H929 cells stably express B cell maturation antigen (BCMA)-a novel treatment target for multiple myeloma (MM) and have been widely used in cancer research and drug development. NCI-H929/GFP cells stably stably express GFP, can be used for *in vitro* assays and *in vivo* imaging.




Figure 1. GFP expression in NCI-H929/GFP stable cell line.

Cell line description

Organism: Homo sapiens (human)

Tissue: Bone Marrow

Cell Type: B lymphocyte

Morphology: Lymphoblast

Culture Properties: Suspension

Disease: Plasmacytoma

Biosafety Level: 2

Medium

1. Complete culture medium: RPMI-1640, 10% fetal bovine serum (FBS), 0.05 mM 2-mercaptoethanol.
1 µg/mL of puromycin may be added to the culture medium. Puromycin should not be added until a culture has been well established from the thawed cells.
2. Freeze medium: Fetal bovine serum (FBS), 6% DMSO

Culture procedure

Thawing of frozen cells

1. Thaw the frozen cryovial by gentle agitation in a 37 °C water bath in 1-2 minutes.
2. Remove the cryovial from the water bath as soon as the contents are thawed, and decontaminate by wiping with 70% ethanol.
3. Transfer the thawed cell suspension to a centrifuge tube containing 10 ml of Complete culture medium, centrifuge at 500 g for 5 minutes.
4. Remove the medium by aspiration, resuspend the cells with 2 ml of the Complete culture medium by gently pipetting up and down.
5. Transfer the cells to a T-25 suspension cell culture flask.
6. Place the cells in a 37°C incubator with 5% CO₂.

Sub-culturing

Cultures can be maintained by the addition of fresh medium or replacement of medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 4×10^5 viable cells/ml. Maintain cell density between 4×10^5 and 1×10^6 viable cells/ml.

Renew or add fresh medium every 2-3 days.